

ACKNOWLEDGMENTS

Stephane Burtey was supported by an ERA-EDTA-EMBO fellowship.

1. Mrug M, Zhou J, Woo Y *et al.* Overexpression of innate immune response genes in a model of recessive polycystic kidney disease. *Kidney Int* 2008; **73**: 63–76.
2. Riera M, Burtey S, Fontes M. Transcriptome analysis of a rat PKD model: Importance of genes involved in extracellular matrix metabolism. *Kidney Int* 2006; **69**: 1558–1563.
3. Werder AA, Amos MA, Nielsen AH *et al.* Comparative effects of germfree and ambient environments on the development of cystic kidney disease in CFWwd mice. *J Lab Clin Med* 1984; **103**: 399–407.
4. Gattone II VH, Cowley Jr BD, Barash BD *et al.* Methylprednisolone retards the progression of inherited polycystic kidney disease in rodents. *Am J Kidney Dis* 1995; **25**: 302–313.

SJ Burtey¹, M Riera¹ and M Fontès¹

¹Department of Medical Genetics and Development, INSERM UMR 491, IPHM, GDR CNRS 2688, Faculté de Médecine de la Timone, Marseille Cedex, France

Correspondence: SJ Burtey, Karsenti Group, Cell Biology and Biophysics Unit, EMBL Heidelberg, Meyerhofstraße, Heidelberg 1 69117, Germany.

E-mail: stephane.burtey@embl.de

Response to ‘Overexpression of complement-component genes in Han:SPRD rats a model of polycystic kidney disease’

Kidney International (2008) **73**, 1325; doi:10.1038/ki.2008.52

We appreciate the letter submitted by Dr Burtey and co-workers in response to our recent report ‘Overexpression of innate immune response genes in a model of recessive polycystic kidney disease’.¹ As noted in their letter,² these investigators observed that, similar to our observations in the *cpk* mouse model, genes encoding multiple innate immune factors, including complement system components, were overexpressed in Han:SPRD-Cy rat kidneys. Furthermore, in reviewing the table in their letter, we note additional gene expression similarities between these two models. For example, in *cpk* kidneys, we also observed increased expression of *Adamts1*, *Ctsd*, *Ctsk*, *Hmox1*, *Lyz* (under alias *Lzp-s*), *Mmp14*, and *Vim*. In addition, among the genes overexpressed in Han:SPRD-Cy kidneys, we have also observed increased expression of *F3*, *Lgals3*, *Mal*, and *Runx1* in *cpk* kidneys (data not shown in our original report). These significant parallels in gene expression profiles indicate that similar immune response pathways are activated in kidneys from two phenotypically distinct rodent models, suggesting that this perturbation may be a common signature of polycystic kidney disease.

1. Mrug M, Zhou J, Woo Y *et al.* Overexpression of innate immune response genes in a model of recessive polycystic kidney disease. *Kidney Int* 2008; **73**: 63–76.
2. Burtey S, Riera M, Fontes M. Overexpression of complement components genes in Han:SPRD rats a model of polycystic kidney disease. *Kidney Int* 2007 (in press).

M Mrug¹, J Zhou¹, Y Woo², X Cui^{1,3}, AJ Szalai^{1,4}, J Novak⁴, GA Churchill² and LM Guay-Woodford^{1,5,6}

¹Department of Medicine, Division of Genetic and Translational Medicine, The University of Alabama at Birmingham, Birmingham, Alabama, USA; ²The Jackson Laboratory, Bar Harbor, Maine, USA; ³Department of Biostatistics, The University of Alabama at Birmingham, Birmingham, Alabama, USA; ⁴Department of Microbiology, The University of Alabama at Birmingham, Birmingham, Alabama, USA; ⁵Department of Genetics, The University of Alabama at Birmingham, Birmingham, Alabama, USA and ⁶Department of Pediatrics, The University of Alabama at Birmingham, Birmingham, Alabama, USA

Correspondence: M Mrug, Department of Medicine, Division of Genetic and Translational Medicine, University of Alabama at Birmingham, 752 Kaul Human Genetics Building, 720 20th Street South, Birmingham, Alabama 35294, USA. E-mail: mmrug@uab.edu or LM Guay-Woodford, Department of Medicine, Division of Genetic and Translational Medicine, University of Alabama at Birmingham, 740 Kaul Human Genetics Building, 720 20th Street South, Birmingham, Alabama 35294, USA. E-mail: lgw@uab.edu

Need for a more realistic cut-off GFR value to define chronic renal failure

Kidney International (2008) **73**, 1325–1326; doi:10.1038/ki.2008.54

To the Editor: We read with interest the article titled ‘Age- and gender-specific reference values of estimated GFR in Caucasians: the Nijmegen Biomedical Study.’ by Wetzels *et al.*,¹ and the commentary by Poggio *et al.*² titled ‘Can we do better than a single estimated GFR threshold when screening for chronic kidney disease?’ published in the September 2007 issue. We have performed a similar work in Indian population where we have measured glomerular filtration rate (GFR) in 610 healthy adults and demonstrated a mean GFR of 81.4 ml min⁻¹ compared with 83–85 ml min⁻¹ reported by Wetzels *et al.*^{1,3}

The definition of decreased GFR relies on an understanding of the ‘normal’ GFR range. The Kidney Disease Outcome Quality Initiative guideline accepts normal GFR as 120 ml min⁻¹ and a 50% reduction in GFR (<60 ml min⁻¹) constitutes chronic renal failure (CRF).⁴ It is universally accepted that GFR declines relentlessly with age; however, Kidney Disease Outcome Quality Initiative guideline assumes that GFR is stable between 18 and 70 years. This assumption allows a single cut-off GFR value for CRF. However, as has been demonstrated by Wetzels *et al.*, a 60 ml min⁻¹ cut-off tends to over-diagnose CRF. This problem gets accentuated in India, where normal GFR is itself lower by 25–30%. GFR of a 20-year-old Indian ranges from 89 to 109 ml min⁻¹ and a 50% reduction from normal translates into 45 ml min⁻¹. We agree with Wetzels *et al.* that GFR of 60 ml min⁻¹ cannot be used to define CRF. Consideration should be given to reduce the cut-off value for defining CRF, and we believe 45 ml min⁻¹ is more realistic in the Indian population.

1. Wetzels JF, Kiemeny LA, Swinkels DW, Willems HL, den Heijer M. Age- and gender-specific reference values of estimated GFR in Caucasians: the Nijmegen Biomedical Study. *Kidney Int* 2007; **72**: 632–637.
2. Poggio ED, Rule AD. Can we do better than a single estimated GFR threshold when screening for chronic kidney disease? *Kidney Int* 2007; **72**: 534–536.

3. Barai S, Bandopadhyaya GP, Patel CD *et al.* Do healthy potential kidney donors in India have an average glomerular filtration rate of 81.4 ml/min? *Nephron Physiol* 2005; **101**: 21–26.
4. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. Kidney Disease Outcome Quality Initiative. *Am J Kidney Dis* 2002; **39**: S1–S266.

S Barai¹ and S Gambhir¹

¹Department of Nuclear Medicine, SGPGIMS, Lucknow, India

Correspondence: S Barai, Department of Nuclear Medicine, SGPGIMS, Raebareli Road, Lucknow 226014, India. E-mail: danzig@rediffmail.com

Response to 'Need for a more realistic cutoff GFR value to define chronic renal failure'

Kidney International (2008) **73**, 1326; doi:10.1038/ki.2008.43

We thank Drs Barai and Gambhir¹ for their thoughtful comments. Indeed, our paper intended to stimulate the discussion on 'normal' values. Obviously, reference values of glomerular filtration rate (GFR) may vary between populations. In the Indian population, GFR appears 20 ml min⁻¹ lower than in the European population.^{2–4} It is important, however, to realize that differences in the methodology of GFR measurement may bias these results. In this respect, we wonder if Barai *et al.*² have used the correction factor of 1.15, which was proposed by Mulligan *et al.*⁵ to improve the accuracy of GFR measurement using the Russel two point method. Thus, a GFR of 100 ml min⁻¹ should be corrected to 115 ml min⁻¹.

Most important is the definition of what is normal. As GFR decreases with age, it seems wise to use age-dependent

reference values as 'normal' values. As such, the use of a fixed GFR cutoff point for defining chronic kidney disease is not logical. Still, the most important step is to define particular cutoff points that are independently associated with an increased risk of morbidity or mortality. Indeed, a recent study suggested that in elderly persons a lower GFR threshold should be used.⁶ Certainly, more epidemiological data in different populations are needed. Meanwhile, we feel that in daily patient care, it is wise to compare the values of an individual patient with values obtained in age- and sex-matched controls.

1. Barai S, Gambhir S. Need for a more realistic cutoff GFR value to define chronic renal failure.
2. Barai S, Bandopadhyaya GP, Pate CD *et al.* Do healthy potential kidney donors in India have an average glomerular filtration rate of 81.4 ml/min. *Nephron Physiol* 2005; **101**: 21–26.
3. Vervoot G, Willems JL, Wetzels JFM. Assessment of glomerular filtration rate in healthy subjects and normoalbuminuric diabetic patients: validation of a new prediction equation. *Nephrol Dial Transplant* 2002; **17**: 1909–1913.
4. Berg UB. Differences in decline in GFR with age between males and females. Reference data on clearances of inulin and PAH in potential kidney donors. *Nephrol Dial Transplant* 2006; **21**: 2577–2582.
5. Mulligan JS, Blue PW, Hasbargen JA. Methods for measuring GFR with Technetium-99m-DTPA: an analysis of several common methods. *J Nucl Med* 1990; **31**: 1211–1219.
6. O'Hare AM, Berthenthal D, Covinsky KE *et al.* Mortality risk stratification in chronic kidney disease: one size for all ages? *J Am Soc Nephrol* 2006; **17**: 846–853.

JFM Wetzels¹

¹Department of Nephrology, Radboud University Nijmegen Medical Center, Nijmegen, The Netherlands

Correspondence: JFM Wetzels, Division of Nephrology, Department of Internal Medicine, Radboud University Nijmegen Medical Centre, PO Box 9101, Nijmegen 6500 HB, The Netherlands. E-mail: j.wetzels@nier.umcn.nl